

Effect of the physical activity program on the treatment of resistant hypertension in primary care

Piotr J. Kruk¹ and Michał Nowicki²

¹Community-Based Health Center in Rząśnia, Rząśnia, Poland and ²Department of Nephrology, Hypertension and Kidney Transplantation, Medical University of Lodz, Lodz, Poland

Research

Cite this article: Kruk PJ and Nowicki M. (2018) Effect of the physical activity program on the treatment of resistant hypertension in primary care. *Primary Health Care Research & Development* 19: 575–583. doi: 10.1017/S1463423618000154

Received: 20 July 2017
Revised: 2 February 2018
Accepted: 11 February 2018
First published online: 22 March 2018

Key words:

ambulatory blood pressure monitoring; physical activity; primary health care; resistant hypertension

Author for correspondence:

Piotr J. Kruk, Community-Based Health Center in Rząśnia, Waryńskiego 6, 98-332 Rząśnia, Poland. E-mail: pikruk@wp.pl

Abstract

Background: Regular physical activity is widely recommended for patients with arterial hypertension as an essential component of lifestyle modification. Much less is known about the impact of physical exercise on the management of treatment of resistant hypertension (RH). The aim was to assess the effect of physical activity program intensified by mobile phone text reminders on blood pressure control in subjects with RH managed in the primary care. **Methods:** In total, 53 patients with primary hypertension were qualified, including 27 who met the criteria for RH and 26 with well-controlled hypertension (WCH). Ambulatory 24-h blood pressure was monitored and body composition evaluated with bioimpedance and habitual physical activity profile was determined continuously over 72 h with accelerometer. All measurements were performed at baseline and after three and six months. The patients were asked to modify their lifestyle according to American Heart Association Guidelines that included regular aerobic physical activity tailored to individual needs. **Findings:** Physical activity in RH increased significantly after six months compared with control subjects ($P=0.001$). Office systolic blood pressure (SBP) and diastolic blood pressure (DBP) in the RH group decreased significantly after three months but after six months only office DBP remained significantly lower. After three months 24-h SBP decreased by 3.1 ± 11 mmHg ($P=0.08$) and DBP by 2.0 ± 6 mmHg ($P=0.17$) in RH, whereas in WCH respective changes were $+1.2 \pm 10$ and -0.3 ± 6 mmHg. After six months 24-h BP changes were similar. **Conclusion:** Individualized structured physical activity program increases physical activity in the treatment of resistant hypertensives in primary care but the effect on 24-h blood pressure is only transient.

Background

Regular physical activity has been recognized as a remedy for many diseases (Kokkinos and Myers, 2010; Booth *et al.*, 2012). Physical activity improves the function of various organs, activates socially, enhances mental well-being, boosts general physical performance, and reduces body mass and mortality (Slentz *et al.*, 2009). The relevant preventive effects of increased physical activity have been studied and confirmed in as many as 35 chronic diseases (Booth *et al.*, 2012).

Individuals with hypertension, in particular those with ineffectively controlled blood pressure, that is with resistant hypertension (RH), have a higher risk of cardiovascular incidents compared with subjects with well-controlled arterial hypertension (WCH) (Calhoun *et al.*, 2008; Mancia *et al.*, 2013).

Treatment of RH remains an important clinical problem despite the advances in pharmacotherapy. RH has been related to 75% greater risk of stroke, 44% greater risk of ischemic heart disease and 30% greater risk of death for any cause as well as an increased risk of heart failure, end-stage renal disease and peripheral artery disease (Muntner *et al.*, 2014). The prevalence of RH reaches 14–16% (Achelrod *et al.*, 2015), and is rising mostly because of the aging of the population and increasing incidence of diabetes and obesity (Roberie and Elliott, 2012). RH is also common in primary care setting and accounts for 9.9–22.8% of all hypertensive patients (Gijón-Conde *et al.*, 2014; Prejbisz *et al.*, 2015).

According to a current definition, RH is diagnosed in patients who, after the administration of antihypertensive drugs from three different groups in therapeutic doses including a diuretic, did not achieve a target blood pressure (BP) $<140/90$ mmHg as well as in those who were administered four and more antihypertensives from three different groups, including a diuretic regardless of achieving a target BP (Calhoun *et al.*, 2008). The population of patients with RH also includes the subjects who do not comply with the doctor's recommendations. It is estimated that as many as 25% of patients may fail to observe systematically the drug regimen (Tomaszewski *et al.*, 2014). The compliance may be even poorer with regard to recommended lifestyle modification, reduction of sodium and alcohol consumption,

introduction of a high-fiber diet, avoiding obesity and increasing physical activity (Mancia *et al.*, 2013; Kraus *et al.*, 2015).

According to several large clinical studies, increased physical activity may have an effect on the reduction of mean BP by up to 4–9 mmHg (Cornelissen and Smart, 2013; Mancia *et al.*, 2013). The benefits of moderate physical activity have also been found in patients with RH (Dimeo *et al.*, 2012), but there were only a few intervention studies carried out in this field. Our initial hypothesis is that the patients with RH are resistant not only to pharmacotherapy, by definition, but also to lifestyle measures including increased level of physical activity.

The aim of this study was to assess the effects of a program of intensified physical activity introduced in primary health care combined with exercise training and short text messages sent to the patients' mobile phones or motivational telephone conversations on BP in patients with treatment of RH.

Methods

The prospective lifestyle intervention study was carried out in a single primary care health center from October 2014 until the end of August 2015. The study protocol was approved by the local Ethics Committee and all the patients gave informed consent before being included. The research was conducted in accordance with the Helsinki Declaration. The participants included a population treated in primary health care in a community-based outpatient clinic in central Poland. The adult population of the Center catchment area is ~3700.

The patients qualified for the study were 18–70 years old and were diagnosed with arterial hypertension according to the 2013 guidelines of the European Society of Hypertension and the European Society of Cardiology (Mancia *et al.*, 2013).

The exclusion criteria included any secondary cause of elevated BP, chronic kidney disease stage 3 and higher, that is estimated glomerular filtration rate (eGFR, 2009 Chronic Kidney Disease – Epidemiology Collaboration, CKD-EPI) <60 mL/min (Levin and Stevens, 2014), mental illness and/or impaired consciousness, significant mobility dysfunction, respiratory diseases, heavy liver failure, active malignant disease, unstable coronary disease or NYHA stage 2–4 heart failure, stroke or transient ischemic attack within the past 12 months.

All patients underwent an extensive diagnostics of RH with accordance to the statement from the American Heart Association (Calhoun *et al.*, 2008) in the reference hypertension center (Department of Nephrology, Hypertension and Kidney Transplantation, Medical University of Lodz) for the secondary causes of arterial hypertension including renal, renovascular and endocrine causes, and pseudohypertension after the diagnosis of the treatment of RH had been established. For safety reasons the subjects were not qualified if they experienced any hypertension urgency or emergency requiring hospitalization for 12 months before the study.

A review of the medical records at the community-based health center allowed identifying patients who initially fulfilled the qualification criteria. After a short visit involving physical examination and analysis of the antihypertensive drug history, and previous diagnostics of arterial hypertension 80 subjects met inclusion criteria and signed an informed consent. After the visit laboratory tests were performed. Eventually 53 patients who fulfilled the clinical and laboratory criteria were enrolled into the study, including 27 who met the criteria for RH (Calhoun *et al.*, 2008) and 26 subjects on antihypertensive medications with well-controlled

BP (Mancia *et al.*, 2013). Most patients who did not meet the criteria had eGFR lower than 60 mL/min. In addition, only the patients who had a long history of the visits in health center, were well known to family doctor and bought the prescribed medications at the pharmacy that is a part of our health center were enrolled. Moreover, the patients were also asked to bring the boxes of used and unused medications at each visit and the pills were counted.

Ambulatory blood pressure monitoring (ABPM) was performed using the Mobil-O-Graph device, version 12 (I.E.M. GmbH, Stolberg, Germany) (Jones *et al.*, 2000). During the day the measurements were taken every 15 min and during the night every 30 min. The time of the day or the night was determined by the patient by pressing the button. We established the following definitions: the circadian dipper pattern as 10–20% reduction of BP measurements at night as compared with ABPM measurements during the day, the mild dipper pattern as 0–10% reduction, the extreme dipper pattern as a reduction of BP measurement by >20%, and the reverse dipper pattern as a rise of BP during the night. Office BP was measured using Accoson Greenlight 300 (A.C. Cossor & Son LTD, Essex, Great Britain).

The analysis of the physical activity profile and energy expenditure was carried out with BodyMedia SenseWear System accelerometer using BodyMedia SenseWear Armband Mini, triaxial accelerometer (BodyMedia Inc., Pittsburgh, PA, USA) attached to the patient's arm for 3 consecutive days (Johannsen *et al.*, 2010).

The body composition analysis using multi-frequency electrical bioimpedance (Nutriguard-M with dedicated Bianostic-AT electrodes; Data Input GmbH, Darmstadt, Germany) was carried out for all patients.

The interview, measurements (ABPM, body composition analysis, physical activity profile) and laboratory tests (complete blood count, creatinine, electrolytes, lipids, glucose, aspartate and alanine transaminases) were repeated after three and six months during the course of the study.

The patients from both groups received the recommendations concerning their diet and healthy lifestyle including physical activity in cardiovascular diseases. The recommendations were based on the American Heart Association's Diet and Lifestyle Recommendations (www.heart.org, 2014) and prepared in writing in Polish language and presented to each patient at the visit and discussed in the form of a short conversation.

At each stage of the study the patients from RH group received also verbal instructions on how to intensify physical activity tailored to their needs and comorbidities. Additionally a meeting with a physical therapist was provided to the patients. During the meeting a demonstration was given how to perform the stretching and the exercises strengthening muscles. Easy-to-apply physical exercises were demonstrated and warm-up elements were discussed. During the 1-h-lasting meeting the patients were predominantly encouraged to engage in physical activity. The aim of this meeting was to decrease an anxiety related to physical exercises known as kinesiphobia (Lundberg *et al.*, 2006). Then the information received was verified during a short conversation. The principles of lifestyle modification were also discussed during all study visits.

After three and six months of the study the patients from RH group received a self-assessment form to be completed at home, which included 10 questions. Questions concerned the difficulties in the implementation exercise, positive or negative attitudes towards physical activity, and the willingness to continue the

exercise (The self-assessment form is available as Online Resource 1). On the basis of the information physical activity was modified and intensified.

In addition, the patients received text messages to their cell phones with reminders about the benefits of regular physical activity three times a week. The messages were tailored to the individual needs of each patient and their content was modified to avoid repeats (Cole-Lewis *et al.*, 2010; de Jongh *et al.*, 2012). Five patients from RH group did not use mobile phones on a daily basis. These patients were encouraged to undertake physical activity during a short landline phone conversation.

The results are presented as percentage values and arithmetic mean with a standard deviation. In order to compare the means of normally distributed variables between groups unpaired *t*-test was used. For non-normally distributed variables Mann-Whitney test was used. The results of repeated measurements were tested with analysis of variance (ANOVA). χ^2 test was used to analyze qualitative variables. For dichotomous variables evaluated repeatedly the Q Cochran test was used. $P < 0.05$ was taken as significant.

Results

Table 1 shows the baseline characteristics of the study groups. No significant differences between groups were observed with regard to age, sex, height, family history of the disease, prevalence of diabetes, ischemic heart disease, chronic kidney disease, alcohol consumption, tobacco smoking, circadian BP patterns or education. Significant differences between groups were found in case of body mass, body mass index (BMI), waist circumference, prevalence of headaches, sleep disturbances including snoring.

Table 2 shows ambulatory and office BP during the study. A significant decrease of office BP was observed in RH group after three months, and the same was true for office diastolic blood pressure (DBP) after six months. There were no statistically significant differences after six months in ABPM values. Systolic blood pressure (SBP) and pulse pressure during the day differed significantly between the groups at baseline but not at the end of the study. During the last three months of the study nighttime ambulatory BP increased significantly in RH group. At the beginning of the study the differences of morning SBP and DBP between groups were numerically noticeable albeit statistically insignificant but that difference was no longer seen at the end of the study.

Figure 1 shows the changes of SBP, DBP and pulse pressure after three and six months of the study compared between patients with RH and WCH. After three months the changes between groups in the 24-h SBP were only borderline significant. No significant difference was demonstrated after six months. However, after excluding five patients who did not use cell phones and receive SMS reminders there was a statistically significant difference between groups in the changes of SBP and pulse pressure after three months ($P = 0.03$ and $P = 0.04$, respectively).

The statistically significant differences in RH were observed in office SBP and DBP (Friedman ANOVA, $P = 0.01$ and $P = 0.02$, respectively), nighttime SBP ($P = 0.02$) and nighttime pulse pressure ($P = 0.002$). No statistically significant differences were found in WCH group.

Figure 2 shows the circadian BP patterns in both groups. No statistically significant differences in the BP patterns were observed in different stages of the study. In RH group a

non-significant reduction of the proportion of patients representing a 'dipper' profile was observed as well as an increase of the 'mild dipper' profile. The total number of the 'dipper' and 'mild dipper' patients increased but the number of patients in the 'extreme dipper' and the 'reverse dipper' category decreased at the end of the study. In WCH group the proportion of 'dippers', 'mild dippers' and 'reverse dippers' decreased.

Table 3 shows the physical activity profiles and the body composition analysis. After six months a statistically significant increase of the number of steps taken and metabolic equivalent and a decrease of both resting and sleeping time were observed in patients with RH. The body composition analysis demonstrated an increase of the total water content and the fat-free body mass in both groups and an increase of the body cell mass between months three and six in RH group ($P = 0.08$). In the first three months of the study a statistically significant increase of extra-cellular mass in RH patients was observed.

In RH patients body mass (89.4 ± 13.6 versus 89.5 ± 14 kg) and BMI (32.5 ± 5.1 versus 32.6 ± 5.3 kg/m²) did not change noticeably while waist circumference decreased (109.5 ± 12 versus 108.6 ± 12 cm; $P = 0.32$). In WCH group, body mass (76.3 ± 12 versus 78 ± 12 kg; $P = 0.07$) and BMI (28.2 ± 4.3 versus 28.8 ± 4.8 kg/m²; $P = 0.05$) increased, whereas the waist circumference did not change (99.3 ± 11 versus 99.4 ± 11 cm).

Mean number of antihypertensive drugs used during the study in RH group slightly decreased from 4.2 ± 0.75 to 3.9 ± 1.2 but the difference was not significant ($P = 0.11$). In WCH patients respective mean values were 1.7 ± 0.5 and 1.8 ± 0.6 ($P = 0.4$).

Median number of drugs was unchanged in both groups in course of the study. At the end of the study the patients with RH completed the self-assessment form. In total, 85% of the patients positively evaluated the program and expressed their willingness to continue the physical activity. In total, 93% declared a subjective improvement of health and nearly all (96%) expressed their readiness to recommend the similar physical activity to their family members.

Discussion

The main finding of our study was that the successful implementation of the individualized structured program of increased physical activity in patients with RH led to moderate and transient BP reduction and to the apparently beneficial changes of body composition.

The study included the patients representing a typical population under the care of a community-based health center. A small size of the local community allowed to closely monitor the course of the study and its progress as well as to prevent non-compliance. The regular visits motivated the patients to intensify physical activity and might have reduced potential concerns related to physical training. Because of the nature of the conversations they were carried out individually but the topics that were discussed were basically the same in each case. The cell phone text messages were also used to encourage the patients to continue and increase the physical activity. This method is quite new but its positive impact of text reminders on the patients' compliance concerning lifestyle changes has already been evaluated in several publications and the method has been gaining popularity (de Jongh *et al.*, 2012; Vodopivec-Jamsec *et al.*, 2012). Bobrow *et al.* found a small but positive effect of transmitting information via SMS to reduce BP in patients with arterial hypertension (Bobrow *et al.*, 2016) but the recent study

Table 1. Baseline characteristics of the patients with resistant and well-controlled hypertension

	Resistant hypertension			Well-controlled hypertension			P value
	All (27)	Women [16 (59%)]	Men [11 (41%)]	All (26)	Women [17 (65%)]	Men [9 (35%)]	
Age (years)	55.5 ± 9	56.5 ± 9	54.2 ± 10	54.8 ± 9	55.7 ± 9	53.1 ± 9	NS
Body weight (kg)	89.4 ± 13.6	85.3 ± 14.7	95.4 ± 9.5	76.3 ± 11.6	73.2 ± 11.8	82.1 ± 9.3	<0.001
Height (m)	1.66 ± 0.1	1.63 ± 0.06	1.7 ± 0.08	1.65 ± 0.1	1.6 ± 0.05	1.72 ± 0.05	NS
BMI (kg/m ²)	32.5 ± 5.1	32.25 ± 5.2	32.9 ± 5.3	28.2 ± 4.3	28.4 ± 4.9	27.7 ± 3.0	<0.005
Waist circumference (cm)	109.5 ± 12.1	107.8 ± 13	111.9 ± 10	99.3 ± 11.1	98.35 ± 12.9	101 ± 6.7	<0.005
Education							
Basic	15 (55.5%)	8 (50%)	7 (63.6%)	11 (42.3%)	8 (47%)	2 (22.2%)	NS
Secondary	9 (33.3%)	6 (37.5%)	3 (27.3%)	11 (42.3%)	5 (29.4%)	6 (66.6%)	
Higher	3 (11.1%)	2 (12.5%)	1 (9.1%)	4 (15.38%)	4 (23.53%)	1 (11.1%)	
Circadian blood pressure pattern							
Dippers	14 (54%)	6 (40%)	8 (73%)	15 (60%)	9 (56%)	6 (67%)	NS
Mild dippers	4 (15%)	3 (20%)	1 (9%)	7 (28%)	4 (25%)	3 (33%)	
Extreme dippers	4 (15%)	3 (20%)	1 (9%)	2 (8%)	2 (13%)	0	
Reverse dippers	4 (15%)	3 (20%)	1 (9%)	1 (4%)	1 (6%)	0	
Coronary heart disease							
Yes	3 (11%)	3 (19%)	0	0	0	0	NS
No	24 (89%)	13 (81%)	11 (100%)	26 (100%)	17 (100%)	9 (100%)	
Diabetes							
Yes	11 (41%)	6 (38%)	5 (45%)	8 (31%)	7 (41%)	1 (11%)	NS
No	16 (59%)	10 (63%)	6 (55%)	18 (69%)	10 (59%)	8 (89%)	
Chronic kidney disease							
Yes	5 (19%)	4 (25%)	1 (9%)	1 (4%)	1 (6%)	0	NS
No	22 (81%)	12 (75%)	10 (91%)	25 (96%)	16 (94%)	9 (100%)	
Chronic back pain							
Yes	17 (63%)	11 (69%)	6 (55%)	10 (38%)	8 (47%)	2 (22%)	0.08
No	10 (37%)	5 (31%)	5 (45%)	16 (62%)	9 (53%)	7 (78%)	
Headache							
Yes	15 (56%)	10 (63%)	5 (45%)	7 (27%)	6 (35%)	1 (11%)	<0.05
No	12 (44%)	6 (38%)	6 (55%)	19 (73%)	11 (65%)	8 (89%)	
Sleep disorders, snoring							
Yes	22 (81%)	14 (88%)	8 (73%)	13 (50%)	8 (47%)	5 (56%)	<0.05
No	5 (19%)	2 (12%)	3 (27%)	13 (50%)	9 (53%)	4 (44%)	
Positive family history of cardiovascular disease							
Yes (hypertension, diabetes, stroke)	25 (93%)	15 (94%)	10 (91%)	22 (85%)	15 (88%)	7 (78%)	NS
No	2 (7%)	1 (6%)	1 (9%)	4 (15%)	2 (12%)	2 (22%)	
Smoking							
Currently, smoking	3 (11%)	1 (6%)	2 (18%)	5 (19%)	3 (18%)	2 (22%)	NS
Non-smoking	23 (85%)	14 (88%)	9 (82%)	18 (69%)	13 (76%)	5 (56%)	
Non-smoker from <5 years ago	1 (4%)	1 (6%)	0	2 (12%)	1 (2%)	2 (22%)	
Alcohol drinking							
Yes	8 (30%)	2 (13%)	6 (55%)	8 (31%)	2 (12%)	6 (67%)	NS
No	19 (70%)	14 (88%)	5 (45%)	18 (69%)	15 (88%)	3 (33%)	

BMI = body mass index.

Data are presented as mean ± SD or number (percentage).

Table 2. 24-h, nighttime, daytime, morning systolic and diastolic blood pressure, pulse pressure in resistant hypertension and well-controlled hypertension groups at baseline and after three and six months of the study

Parameter (mmHg)	Time of measurement	Resistant	Hypertension	Well-controlled hypertension	P value
Office SBP	At baseline	150 ± 24		132 ± 8.5	<0.001
	After 3 months	137.5 ± 17	P = 0.004*	129.7 ± 10	0.1
	After 6 months	140 ± 24	P = 0.08	130.7 ± 11	0.2
Office DBP	At baseline	90 ± 14		80.5 ± 7.5	<0.005
	After 3 months	83 ± 11	P = 0.001*	77.7 ± 7	0.1
	After 6 months	84.4 ± 13	P = 0.04**	80.7 ± 8	0.2
SBP ABPM	At baseline	127 ± 17		119 ± 12	<0.05
	After 3 months	126 ± 15	P = 0.20	120 ± 11	0.1
	After 6 months	127 ± 18	P = 0.60	119 ± 9	0.1
DBP ABPM	At baseline	75.4 ± 11		72.6 ± 7.1	0.3
	After 3 months	73.8 ± 11	P = 0.10	71.9 ± 6.5	0.5
	After 6 months	74.6 ± 12	P = 0.40	71.9 ± 5.2	0.4
Daytime SBP ABPM	At baseline	131.3 ± 16		122.4 ± 12	<0.05
	After 3 months	131.4 ± 14	P = 0.30	124 ± 12	0.1
	After 6 months	130 ± 16	P = 0.30	123 ± 10	0.1
Daytime DBP ABPM	At baseline	77.8 ± 11		75 ± 7	0.3
	After 3 months	77.5 ± 10	P = 0.30	74.7 ± 7	0.3
	After 6 months	76.2 ± 11	P = 0.20	74 ± 5	0.4
Nighttime SBP ABPM	At baseline	116.7 ± 20		109.5 ± 15	0.2
	After 3 months	111.6 ± 17	P = 0.07	107 ± 12	0.3
	After 6 months	119 ± 25	P = 0.70	107 ± 9	0.05
Nighttime DBP ABPM	At baseline	67.4 ± 13		64.8 ± 9	0.8
	After 3 months	65.7 ± 12	P = 0.40	63 ± 6	0.4
	After 6 months	69 ± 15	P = 0.90	64 ± 6	0.2
Pulse pressure ABPM	At baseline	53.2 ± 11		46.4 ± 9	<0.05
	After 3 months	52.3 ± 8	P = 0.20	48 ± 9	0.1
	After 6 months	52.5 ± 10	P = 0.40	47 ± 9	0.1
Nighttime pulse pressure ABPM	At baseline	49 ± 11		44.2 ± 9	0.4
	After 3 months	46 ± 9	P = 0.03**	44 ± 9	0.4
	After 6 months	50 ± 14	P = 0.80	43.5 ± 8	0.1
Daytime pulse pressure ABPM	At baseline	53.5 ± 10		47.6 ± 10	<0.05
	After 3 months	54.7 ± 9	P = 0.98	49.5 ± 10	0.06
	After 6 months	53 ± 9	P = 0.50	49 ± 9	0.1
Morning SBP ABPM	At baseline	136.6 ± 25		125 ± 19	0.06
	After 3 months	136 ± 16	P = 0.40	125 ± 20	0.05
	After 6 months	116.6 ± 37	P = 0.07	120 ± 17	0.3
Morning DBP ABPM	At baseline	80.5 ± 15		73.2 ± 10	0.05
	After 3 months	80.3 ± 14	P = 0.50	74 ± 12	0.1
	After 6 months	75.2 ± 19	P = 0.15	74 ± 12	0.3

ABPM = automatic blood pressure monitoring; BP = blood pressure; DBP = diastolic blood pressure; Morning SBP and DBP = blood pressure mean during 1st hour after waking up; SBP = systolic blood pressure.

Data are presented as mean ± SD.

*P < 0.005 for the difference in the group of resistant hypertension versus baseline value.

**P < 0.05 for the difference in the group of resistant hypertension versus baseline value.




Figure 1. Systolic blood pressure, pulse pressure in resistant hypertension and well-controlled hypertension groups in observation period. Δ = Change of parameter in observation period; DBP = diastolic blood pressure; PP = pulse pressure; SBP = systolic blood pressure. Data are presented as mean; P value of <0.05 is considered statistically significant




Figure 2. Circadian blood pressure patterns during the study. RH = resistant hypertension group; WCH = well-controlled hypertension group

comparing a standard behavioral weight loss intervention versus a technology-enhanced weight loss intervention did not result in greater weight loss (Jakicic *et al.*, 2016).

A study that referred to a similar population demonstrated that a 12-week aerobic exercise program was effective in decreasing BP. The BP reduction achieved during the day was 6 ± 12 and 3 ± 7 mmHg for SBP and DBP, respectively (Dimeo *et al.*, 2012). In our population the respective changes of BP were smaller and non-significant reaching 1.5 ± 16 mmHg and 1.6 ± 11 mmHg. The smaller effect in our study may be explained by the lower 24-h mean SBP and DBP at baseline (127.2 and 75.4 mmHg, respectively) compared with 135.3 and 75.4 mmHg in a study of Dimeo *et al.*

We were not able to offer group activities and the exercises were not supervised or led by a coach for the whole duration of the study. Instead our simplified protocol included the methods that were widely available, inexpensive and that could be implemented without any additional preparation/training. Such program is more likely to be accepted by the patients who are generally reluctant to participate in group activities (Freene *et al.*, 2014). The effectiveness of various methods aimed at increasing the physical activity was analyzed in Australia and that study

confirmed the aptness relevance of our assumptions (Freene *et al.*, 2014; 2015).

Dimeo *et al.* applied 30-min exercises such as walking on a treadmill with a gradual extension of the training duration, three times a week (Dimeo *et al.*, 2012). Guimaraes *et al.* presented a slightly different approach implementing physical activity in the form of strength exercises based on using the persons own body mass (calisthenics) and walking in the water heated up to the temperature of 30–32°C. After 12 weeks of exercises three times a week, including 1-h meeting the lowering effects on BP were observed both during the day and night BP (Guimaraes *et al.*, 2014). Ribeiro *et al.* found a significant effect of physical exercise on BP in individuals with high BP values at the start (Ribeiro *et al.*, 2015). In our study a reference group included the individuals with well-controlled BP using ≤ 3 antihypertensives while in the aforementioned studies control patients had a RH with much higher BP at baseline. We considered a comparative group of well-controlled hypertensive patients with significantly better cardiovascular prognosis as a good background for demonstrating beneficial changes in RH group hence our study showed that the impact of the physical activity program is not inferior in patients with RH compared with the patients with well-controlled hypertension.

A nighttime BP during sleep has been established as an important prognostic factor in recent years, more important than 24-h mean values or morning BP (Salles *et al.*, 2008; Hermida *et al.*, 2013; 2014). Ling *et al.* postulated a beneficial influence of a regular aerobic activity on strengthening the effect of the SBP and pulse pressure drop at night ('dipping') despite the lack of the influence upon the 24-h SBP (Ling *et al.*, 2014). Our study showed also the reduction of the percentage of the patients with the 'dipper' profile and an increase of the number of patients with the 'mild dipper' profile at the end of the study. It needs to be emphasized that the mean nighttime values of BP remained within the normal range during the whole study and at the end the proportion of the 'reverse dipper' and 'extreme dipper' patients known to have the highest risk of cardiovascular incidents (Kario and Shimada, 2004) was lower. The morning SBP tended to decrease [but the difference did not reach the level of statistical significance ($P=0.07$)] in RH group, which might be related to the change of the 24-h pressure profile into a more beneficial pattern as suggested by some authors (Xu *et al.*, 2012).

The importance of diabetes and obesity in the pathogenesis of arterial hypertension and its resistance to treatment is well documented (Calhoun *et al.*, 2008; Roberie and Elliott, 2012). Unfortunately RH group was too small to analyze the effect of obesity and diabetes on the effects of physical activity on the control of hypertension.

The study has other limitations. First, the impact of placebo and Hawthorne effect, that is the knowledge that the person participates in the study upon the results cannot be excluded. The patients were qualified to RH group on the basis of the current definition of RH based on office measurements (Calhoun *et al.*, 2008), and therefore the group might have also included the patients experiencing the apparent RH, such as 'white coat syndrome' and medication non-adherence (Townsend and Epstein, 2016). Importantly the results were primarily analyzed with respect to ABPM that is as the best method for this population (Calhoun *et al.*, 2008; Syrseloudis *et al.*, 2011; Mancia *et al.*, 2013).

The lower reduction of BP than in other studies might have been a consequence of a less intensive exercise program.

Table 3. Physical activity, energy expenditure and body composition in patients with resistant hypertension and well-controlled hypertension

Parameter	Time of measurement	Resistant Hypertension	Well-controlled hypertension	P value
Physical activity of moderate or higher intensity, min	At baseline	290 ± 254	373 ± 220	0.06
	After 3 months	273 ± 144	424 ± 293	<i>P</i> = 0.97
	After 6 months	363 ± 284	421 ± 279	<i>P</i> = 0.30
Number of steps taken during 3 days	At baseline	17 361 ± 6815	21 374 ± 9481	0.08
	After 3 months	20 807 ± 8539	243 314 ± 11 135	<i>P</i> = 0.05
	After 6 months	23067 ± 7741	25 779 ± 11 201	<i>P</i> = 0.002*
MET average, during 3 days	At baseline	1.325 ± 0.3	1.48 ± 0.2	<0.05
	After 3 months	1.33 ± 0.2	1.52 ± 0.3	<i>P</i> = 0.80
	After 6 months	1.46 ± 0.3	1.55 ± 0.3	<i>P</i> = 0.001*
Energy expenditure associated with physical activity during 3 days, (J)	At baseline	6850 ± 5812	7585 ± 4708	0.27
	After 3 months	6500 ± 3303	8861 ± 6505	<i>P</i> = 0.90
	After 6 months	8327 ± 5436	8507 ± 6098	<i>P</i> = 0.20
Total energy expenditure during 3 days (J)	At baseline	34 950 ± 6193	33 054 ± 6753	0.15
	After 3 months	35 310 ± 5633	34 426 ± 9123	<i>P</i> = 0.90
	After 6 months	37 769 ± 7326	34 172 ± 8048	<i>P</i> = 0.06
Rest time in 3 days, including a sedentary lifestyle (min)	at baseline	1573 ± 263	1520 ± 175	0.24
	After 3 months	1511 ± 263	1573 ± 312	<i>P</i> = 0.30
	After 6 months	1458 ± 292	1483 ± 246	<i>P</i> = 0.03**
Sleep during 3 days (min)	At baseline	1278 ± 228	1298 ± 156	0.72
	After 3 months	1238 ± 246	1293 ± 218	<i>P</i> = 0.80
	After 6 months	1147 ± 270	1281 ± 212	<i>P</i> = 0.02**
Total body water (l)	At baseline	43 ± 7.2	39.1 ± 6.6	<0.05
	After 3 months	43.6 ± 7.4	39.1 ± 7.3	<i>P</i> = 0.08
	After 6 months	44 ± 7.3	40.3 ± 6.9	<i>P</i> = 0.01**
Lean body mass (kg)	At baseline	58.8 ± 9.8	53.4 ± 9.1	<0.05
	After 3 months	59.6 ± 10.2	53.4 ± 10	<i>P</i> = 0.11
	After 6 months	60.1 ± 9.9	55.1 ± 9.4	<i>P</i> = 0.01**
Fat mass (kg)	At baseline	29.9 ± 9.1	22.6 ± 8.1	<0.005
	After 3 months	30.6 ± 9.6	23.8 ± 8.1	<i>P</i> = 0.06
	After 6 months	30.1 ± 9.5	23.3 ± 9.1	<i>P</i> = 0.77
Extracellular mass (kg)	At baseline	27.8 ± 4.7	25.3 ± 4.0	<0.05
	After 3 months	28.9 ± 4.5	25.4 ± 4.4	<i>P</i> = 0.005*
	After 6 months	28.4 ± 5.5	26.1 ± 4.4	<i>P</i> = 0.36
Body cell mass (kg)	At baseline	30.7 ± 6.5	28.1 ± 5.6	0.12
	After 3 months	30.6 ± 6.4	28 ± 6.2	<i>P</i> = 0.84
	After 6 months	31.7 ± 6.1	29 ± 5.6	<i>P</i> = 0.10
Basal metabolic rate (kcal)	At baseline	1586 ± 26	1502 ± 177	0.12
	After 3 months	1584 ± 23	1500 ± 194	<i>P</i> = 0.86
	After 6 months	1616 ± 195	1531 ± 178	<i>P</i> = 0.10

MET = metabolic equivalent, 1 MET responsible resting oxygen consumption.

Data are presented as mean ± SD; *Conversion factors to SI units are as follows: for basal metabolic rate, 4184; for total body water, 0.001'.

**P* < 0.005 for the difference in the group of resistant hypertension versus baseline value.

***P* < 0.05 for the difference in the group of resistant hypertension versus baseline value.

In addition, the strongest effect on BP reduction was observed in the groups treated with isometric resistance exercises but that type of training was not used in our study (Cornelissen and Smart, 2013).

The strength of the study is the intervention that is simple, inexpensive and accessible in primary care. It is based on a tailor-made approach and provides an opportunity to select the most suitable type of activity. It is also well tolerated and well assessed by patients.

In conclusion the regular physical activity is a multidirectional form of intervention that has a beneficial impact on the patients with hypertension. Its benefits in patients with treatment of RH are however moderate and limited in time. It seems commendable to try to physically activate patients with RH in addition to a pharmacological therapy.

Acknowledgments. The authors thank the patients without whom this study would not have been possible. They also especially thank the staff of the Community-Based Health Center in Rzażnia and the staff of the Department of Nephrology, Hypertension and Kidney Transplantation in Lodz for their support.

Financial Support. This work was supported by the Medical University of Lodz, Poland (M.N., grant number 503/1-151-02/503-0).

Conflicts of Interest. The authors declare no conflicts of interest to disclose.

Ethical Standards. The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional guidelines on human experimentation (Bioethics Committee of the Medical University of Lodz) and with the Helsinki Declaration of 1975, as revised in 2008. Bioethics Committee of the Medical University of Lodz approved this project (Resolution No. RNN/169/14/14/EC of 10.28.2014).

References

- Achelrod D, Wenzel U and Frey S** (2015) Systematic review and meta-analysis of the prevalence of resistant hypertension in treated hypertensive populations. *American Journal of Hypertension* **28**, 355–361.
- American Heart Association.** The American Heart Association's diet and lifestyle recommendations. Retrieved 25 October 2014 from <https://www.heart.org/>.
- Bobrow K, Farmer AJ, Springer D, Shannyinde M, Yu LM, Brennan T, Rayner B, Namane M, Steyn K, Tarassenko L and Levitt N.** (2016) Mobile phone text messages to support treatment adherence in adults with high blood pressure (SMS-Text Adherence Support [StAR]): a single-blind, randomized trial. *Circulation* **133**, 592–600.
- Booth FW, Roberts CK and Laye MJ** (2012) Lack of exercise is a major cause of chronic diseases. *Comprehensive Physiology* **2**, 1143–1211.
- Calhoun DA, Jones D, Textor S, Goff DC, Murphy TP, Toto RD, White A, Cushman WC, White W, Sica D, Ferdinand K, Giles TD, Falkner B and Carey RM** (2008) Resistant hypertension: diagnosis, evaluation, and treatment. A scientific statement from the American Heart Association Professional Education Committee of the Council for High Blood Pressure Research. *Hypertension* **51**, 1403–1419.
- Cole-Lewis H and Kershaw T** (2010) Text messaging as a tool for behavior change in disease prevention and management. *Epidemiologic Reviews* **32**, 56–69.
- Cornelissen VA and Smart NA** (2013) Exercise training for blood pressure: a systematic review and meta-analysis. *Journal of the American Heart Association* **2**, e004473.
- de Jongh T, Gurol-Urganci I, Vodopivec-Jamsec V, Car J and Atun R** (2012) Mobile phone messaging for facilitating self-management of long-term illnesses. *Cochrane Database of Systematic Reviews* **12**, CD007459.
- Dimeo F, Pagonas N, Seibert F, Arndt R, Zidek W and Westhoff TH** (2012) Aerobic exercise reduces blood pressure in resistant hypertension. *Hypertension* **60**, 653–658.
- Freene N, Waddington G, Chesworth W, Davey R and Cochrane T** (2014) Community group exercise versus physiotherapist-led home-based physical activity program: barriers, enablers, and preferences in middle-aged adults. *Physiotherapy Theory and Practice* **30**, 85–93.
- Freene N, Waddington G, Davey R and Cochrane T** (2015) Longitudinal comparison of a physiotherapist-led, home-based and group-based program for increasing physical activity in community-dwelling middle-aged adults. *Australian Journal of Primary Health* **21**, 189–196.
- Gijón-Conde T, Graciani A and Banegas JR** (2014) Resistant hypertension: demography and clinical characteristics in 6,292 patients in a primary health care setting. *Revista Española de Cardiología (Engl Ed)* **67** (4), 270–276.
- Guimaraes GV, de Barros Cruz LG, Fernandes-Silva MM, Dorea EL and Bocchi EA** (2014) Heated water-based exercise training reduces 24-hour ambulatory blood pressure levels in resistant hypertensive patients: a randomized controlled trial (HEX trial). *International Journal of Cardiology* **172**, 434–441.
- Hermida RC, Ayala DE, Fernández JR and Mojón A** (2013) Sleep-time blood pressure: prognostic value and relevance as a therapeutic target for cardiovascular risk reduction. *Chronobiology International* **30**, 68–86.
- Hermida RC, Ayala DE, Mojón A, Smolensky MH, Portaluppi F and Fernández JR** (2014) Sleep-time ambulatory blood pressure as a novel therapeutic target for cardiovascular risk reduction. *Journal of Human Hypertension* **28**, 567–574.
- Jakicic JM, Davis KK, Rogers RJ, King WC, Marcus MD, Helsel D, Rickman AD, Wahed AS and Belle SH.** (2016) Effect of wearable technology combined with a lifestyle intervention on long-term weight loss: the IDEA randomized clinical trial. *JAMA* **316**, 1161–1171.
- Johannsen DL, Calabro MA, Stewart J, Franke W, Rood JC and Welk GJ** (2010) Accuracy of armband monitors for measuring daily energy expenditure in healthy adults. *Medicine and Science in Sports and Exercise* **42**, 2134–2140.
- Jones C, Taylor K, Chowienczyk P, Poston L and Shennan AH** (2000) A validation of the Mobil O Graph (version 12) ambulatory blood pressure monitor. *Blood Pressure Monitoring* **5**, 233–238.
- Kario K and Shimada K** (2004) Risers and extreme-dippers of nocturnal blood pressure in hypertension: antihypertensive strategy for nocturnal blood pressure. *Clinical and Experimental Hypertension* **26**, 177–189.
- Kokkinos P and Myers J** (2010) Exercise and physical activity: clinical outcomes and applications. *Circulation* **122**, 1637–1648.
- Kraus WE, Bittner V, Appel L, Blair SN, Church T, Després JP, Franklin BA, Miller TD, Pate RR, Taylor-Piliae RE, Vafiadis DK and Whitte L.** (2015) The national physical activity plan: a call to action from the American Heart Association: a science advisory from the American Heart Association. *Circulation* **131**, 1932–1940.
- Levin A and Stevens P** (2014) Summary of KDIGO 2012 CKD Guideline. Behind the scenes, need for guidance, and a framework for moving forward. *Kidney International* **85**, 49–61.
- Ling C, Diaz KM, Kretzschmar J, Fearheller DL, Sturgeon KM, Perkins A, Veerabhadrapa P, Williamson ST, Lee H, Grimm H, Babbitt DM and Brown MD** (2014) Chronic aerobic exercise improves blood pressure dipping status in African American nondippers. *Blood Pressure Monitoring* **19**, 353–358.
- Lundberg M, Larsson M, Östlund H and Styf J** (2006) Kinesiophobia among patients with musculoskeletal pain in primary healthcare. *Journal of Rehabilitation Medicine* **38**, 37–43.
- Mancia G, Fagard R, Narkiewicz K, Redón J, Zanchetti A, Böhm M, Christiaens T, Cifkova R, De Backer G, Dominiczak A, Galderisi M, Grobbee DE, Jaarsma T, Kirchhof P, Kjeldsen SE, Laurent S, Manolis AJ, Nilsson PM, Ruilope RM, Schmieder RE, Sirnes PA, Sleight P, Viigimaa M, Waeber B and Zannad F** (2013) 2013ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Journal of Hypertension* **31**, 1281–1357.
- Muntner P, Davis BR, Cushman WC, Bangalore S, Calhoun DA, Pressel SE, Black HR, Kostis JB, Probstfield JL, Whelton PK and Rahman M** (2014) Treatment – resistant hypertension and the incidence of cardiovascular disease and end-stage renal disease: results from the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *Hypertension* **64**, 1012–1021.

- Prejbisz A, Klocek M, Gašowski J, Topór-Mądry R, Leśniak W, Kabat M, Czarnecka D, Kawecka-Jaszcz K, Narkiewicz K and Januszewicz A** (2015) Factors associated with resistant hypertension in a large cohort of hypertensive patients: the Pol-Fokus study. *Polskie Archiwum Medycyny Wewnętrznej* **125**, 249–259.
- Ribeiro F, Costa R and Mesquita-Bastos J** (2015) Exercise training in the management of patients with resistant hypertension. *World Journal of Cardiology* **7**, 47–51.
- Roberie DR and Elliott WJ** (2012) What is the prevalence of resistant hypertension in the United States? *Current Opinion in Cardiology* **27**, 386–391.
- Salles GF, Cardoso CR and Muxfeldt ES** (2008) Prognostic influence of office and ambulatory blood pressures in resistant hypertension. *Archives of Internal Medicine* **168**, 2340–2346.
- Slentz CA, Houmard JA and Kraus WE** (2009) Exercise, abdominal obesity, skeletal muscle, and metabolic risk: evidence for a dose response. *Obesity* **17**, 27–23.
- Syrseoudis D, Andrikou I, Andrikou E, Dimitriadis K and Stefanadis C** (2011) Ambulatory blood pressure monitoring in resistant hypertension. *International Journal of Hypertension* **2011**, 285612.
- Tomaszewski M, White C, Patel P, Masca N, Damani R, Hepworth J, Samani NL, Gupta P, Madira W, Stanley A and Williams B** (2014) High rates of non-adherence to antihypertensive treatment revealed by high-performance liquid chromatography-tandem mass spectrometry (HP LC-MS/MS) urine analysis. *Heart* **100**, 855–861.
- Townsend RR and Epstein M** (2016) Resistant hypertension: insights on evaluation and management in the Post-SPRINT (systolic blood pressure intervention trial) era. *Hypertension* **68**, 1073–1080.
- Vodopivec-Jamsec V, de Jongh T, Gurol-Urganci I, Atun R and Car J** (2012) Mobile phone messaging for preventive health care. *Cochrane Database Systematic Reviews* **12**, CD007457.
- Xu T, Zhang YQ and Tan XR** (2012) The dilemma of nocturnal blood pressure. *Journal of Clinical Hypertension* **14**, 787–791.